Neuroscience Letters, 147 (1992) 237–239
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Agranoff, B.W., see Lyon, M.J., 72
Akamatsu, N., see Inenaga, K., 189
Åkerman, K.E.O. and Grönblad, M., Intracellular free [Ca^{2+}] and [Na^{+}] in response to capsaicin in cultured dorsal root ganglion cells, 13
Allen, S., see Griffiths, R., 175
Aloe, L., see Laudiero, L.B., 9
Anderton, B.H., see Lantos, P.L., 67
Anderton, B.H., see Hanger, D.P., 58
Asano, T., see Nagafuji, T., 159
Atlas, D., see Osherov, A., 233

Baker, H., see Kang, U.J., 53
Barbeito, L., see Chéramy, A., 209
Bargas, J., see García-Ugalde, G., 229
Behrens, U.D., Wagner, H.-J. and Kirsch, M., cAMP-mediated second messenger mechanisms are involved in spinule formation in teleost cone horizontal cells, 93
Bennett, M.R., see Karunanithi, S., 49
Bentivoglio, M., see Chen, S., 85
Bloms, P., Mußhoff, U., Madeja, M., Müsch-Nittel, K., Kuhlmann, D., Spener, F. and Speckmann, E.-J., Suppression of a ligand operated membrane current by the epileptogenic agent pentylenetetrazol in oocytes of *Xenopus laevis* after injection of rat brain RNA, 155
Bloms, P., see Mußhoff, U., 163
Bouisset, S., see Nouillot, P., 1
Brailowsky, S., see García-Ugalde, G., 229
Breidbach, O., Dennis, R., Marx, J., Görlach, C., Wiegandt, H. and Wegerhoff, R., Insect glial cells show differential expression of a glycolipid-derived, glucuronic acid-containing epitope throughout neurogenesis: detection during postembryogenesis and regeneration in the central nervous system of *Tenebrio molitor* L., 5
Brion, J.-P., see Hanger, D.P., 58
Brookes, N., Regulation of the glutamine content of astrocytes by cAMP and hydrocortisone: effect of pH, 139
Bruce, M.E., see Jeffrey, M., 106
Bruce, M.T., see Lantos, P.L., 67
Buttinelli, C., see Laudiero, L.B., 9

Champagnat, J., see Fortin, G., 89
Chéramy, A., Barbeito, L., Godeheu, G. and Glowinski, J., Riluzole inhibits the release of glutamate in the caudate nucleus of the cat in vivo, 209
Chen, S., Raos, V. and Bentivoglio, M., Connections of the thalamic reticular nucleus with the contralateral thalamus in the rat, 85
Chiadò-Piat, L., see Mongini, T., 37
Choi, C.R., see Park, C.K., 41
Clinton, J., see Lantos, P.L., 67
Collinge, J., see Lantos, P.L., 67
Copenhagen, D.R., see Dixon, D.B., 147
Corasaniti, M.T., Tartaglia, R.L., Melino, G., Nisticò, G. and Finazzi-Agrò, A., Evidence that CHP100 neuroblastoma cell death induced by *N*-methyl-D-aspartate involves L-arginine-nitric oxide pathway activation, 221
Crow, T., see Falk-Vairant, J., 45

Denavit-Saubié, M., see Fortin, G., 89
Dennis, R., see Breidbach, O., 5
Dixon, D.B. and Copenhagen, D.R., Suppression of a calcium current by CNQX and kynurenate, 147
Djamgoz, M.B.A., see Ter-Margarian, A., 131
Do, M.C., see Nouillot, P., 1
Dodt, H.-U., see Pawelzik, H., 167
Doey, L.J., see Lantos, P.L., 67
Domingo, J., see Ferriero, D.M., 29
Donevan, S.D., see Subramaniam, S., 213
Doriguzzi, C., see Mongini, T., 37
Du, F. and Schwarcz, R., Aminooxyacetic acid causes selective neuronal loss in layer III of the rat medial entorhinal cortex, 185
Dun, N.J., Dun, S.L., Forstermann, U. and Tseng, L.F., Nitric oxide synthase immunoreactivity in rat spinal cord, 217
Dun, S.L., see Dun, N.J., 217

Evercooren, A.B.-V., see Gansmuller, A., 151

Falk-Vairant, J. and Crow, T., Intracellular injections of BAPTA block induction of enhancement in *Hermissenda* type B-photoreceptors, 45
Ferriero, D.M., Sheldon, R.A. and Domingo, J., Somatostatin is altered in developing retina from ethanol-exposed rats, 29
Finazzi-Agrò, A., see Corasaniti, M.T., 221
Forstermann, U., see Dun, N.J., 217
Fortin, G., Velluti, J.C., Denavit-Saubié, M. and Champagnat, J., Responses to repetitive afferent activity of rat solitary complex neurons isolated in brainstem slices, 89

Galarraga, E., see García-Ugalde, G., 229
Gansmuller, A., Kruger, F., Gumpel, M. and Evercooren, A.B.-V., Photoconverted carbocyanine DiI allows direct visualization of transplanted glial cells at the ultrastructural level, 151
García-Ugalde, G., Galarraga, E., Bargas, J. and Brailowsky, S., Hyperexcitability of hippocampal CA1 region in brain slices after GABA withdrawal, 229
Garfield, R.E., see Khan, I., 115
Gebicke-Haerter, P.J., see Nörenberg, W., 171
Gillessen, S., see Laudiero, L.B., 9
Glowinski, J., see Chéramy, A., 209
Godeheu, G., see Chéramy, A., 209
Goji, Y., see Yamamoto, H., 97
Goldberg, D., see Osherov, A., 233
Goodsir, C.M., see Jeffrey, M., 106
Görlach, C., see Breidbach, O., 5
Grieve, A., see Griffiths, R., 175
Griffiths, R., Grieve, A., Allen, S. and Olverman, H.J., Neuronal and glial plasma membrane carrier-mediated uptake of L-homocysteate is not selectively blocked by β-*p*-chlorophenylglutamate, 175
Grönblad, M., see Åkerman, K.E.O., 13
Grover, A.K., see Khan, I., 115
Gumpel, M., see Gansmuller, A., 151

Halliday, W.G., see Jeffrey, M., 106
Hanger, D.P., Hughes, K., Woodgett, J.R., Brion, J.-P. and Anderton, B.H., Glycogen synthase kinase-3 induces Alzheimer's disease-like phosphorylation of tau: generation of paired helical filament epitopes and neuronal localisation of the kinase, 58
Heaton, M.B., see Walker, D.W., 77
Héron, A., Lasbennes, F. and Seylaz, J., Effect of two different routes

of administration of *R*-PIA on glutamate release during ischemia, 205
Honma, K.-i., see Honma, S., 201
Honma, S., Kanematsu, N., Katsuno, Y. and Honma, K.-i., Light suppression of nocturnal pineal and plasma melatonin in rats depends on wavelength and time of day, 201
Hori, K., see Tanaka, J., 111
Hua, H.T., see McNeill, D.L., 33
Hughes, K., see Hanger, D.P., 58
Hulsebosch, C.E., see Stroemer, R.P., 21
Hunter, B.E., see Walker, D.W., 77

Illes, P., see Nörenberg, W., 171
Inagaki, M., see Kiuchi, Y., 193
Inenaga, K., Akamatsu, N., Nagatomo, T., Ueta, Y. and Yamashita, H., Intracellular EGTA alters phasic firing of neurons in the rat supraoptic nucleus in vitro, 189
Ito, T., see Komori, K., 197
Izumi, J., see Kiuchi, Y., 193

Janota, I., see Lantos, P.L., 67
Jeffrey, M., Goodsir, C.M., Bruce, M.E., McBride, P.A., Scott, J.R. and Halliday, W.G., Infection specific prion protein (PrP) accumulates on neuronal plasmalemma in scrapie infected mice, 106
Joh, T.H., see Kang, U.J., 53
Jones, S.V.P., m4 muscarinic receptor subtype activates an inwardly rectifying potassium conductance in AtT20 cells, 125

Kane, J.M., see Kehne, J.H., 101
Kanematsu, N., see Honma, S., 201
Kang, J.K., see Park, C.K., 41
Kang, U.J., Park, D.H., Wessel, T., Baker, H. and Joh, T.H., DOPA-decarboxylation in the striata of rats with unilateral substantia nigra lesions, 53
Karunanithi, S., Lavidis, N.A. and Bennett, M.R., The effect of adenosine on spontaneous and evoked quantal secretion from different release sites of amphibian motor-nerve terminals, 49
Kasahara, Y., see Komori, K., 197
Katsumaru, H., see Saito, Y., 81
Katsuno, Y., see Honma, S., 201
Kehne, J.H., Ketteler, H.J., Kane, J.M., McCloskey, T.C., Senyah, Y. and Palfreyman, M.G., MDL 27,531 reduces spontaneous hindlimb contractions in rats with chronic transections of the spinal cord, 101
Kent, T.A., see Stroemer, R.P., 21
Ketteler, H.J., see Kehne, J.H., 101
Khan, I., Tabb, T., Garfield, R.E. and Grover, A.K., Polymerase chain reaction assay of mRNA using 28S rRNA as internal standard, 115
King, M.A., see Walker, D.W., 77
Kirsch, M., see Behrens, U.D., 93
Kiuchi, Y., Inagaki, M., Izumi, J., Matsumoto, M., Yamazaki, Y. and Oguchi, K., Effect of local cyanide perfusion on rat striatal extracellular dopamine and its metabolites as studied by in vivo brain microdialysis, 193
Koide, T., see Nagafuji, T., 159
Koito, H., see Tanaka, J., 111
Komori, K., Kunimi, Y., Yamaoka, K., Ito, T., Kasahara, Y. and Nagatsu, I., Semiquantitative analysis of immunoreactivities of tyrosine hydroxylase and aromatic L-amino acid decarboxylase in the locus coeruleus of desipramine-treated mice, 197
Kruger, F., see Gansmuller, A., 151
Kuhlmann, D., see Bloms, P., 155
Kunimi, Y., see Komori, K., 197

Lantos, P.L., McGill, I.S., Janota, I., Doey, L.J., Collinge, J., Bruce, M.T., Whatley, S.A., Anderton, B.H., Clinton, J., Roberts, G.W. and Rossor, M.N., Prion protein immunocytochemistry helps to establish the true incidence of prion diseases, 67
Lasbennes, F., see Héron, A., 205
Laudiero, L.B., Aloe, L., Levi-Montalcini, R., Buttinelli, C., Schilter, D., Gillessen, S. and Otten, U., Multiple sclerosis patients express increased levels of β-nerve growth factor in cerebrospinal fluid, 9
Lavidis, N.A., see Karunanithi, S., 49
Lee, N., see Walker, D.W., 77
Levi-Montalcini, R., see Laudiero, L.B., 9
Lyon, M.J., Agranoff, B.W., Sokoloff, L. and Smith, C.B., Residual effects of tracer in sequential double label deoxyglucose studies, 72

Madeja, M., see Bloms, P., 155
Madeja, M., see Mußhoff, U., 163
Mailleux, P., Zhang, F. and Vanderhaeghen, J.-J., The dopamine D_1 receptor antagonist SCH-23390 decreases the mRNA levels of the transcription factor zif268 (krox-24) in adult rat intact striatum – an in situ hybridization study, 182
Mailleux, P. and Vanderhaeghen, J.-J., Age-related loss of cannabinoid receptor binding sites and mRNA in the rat striatum, 179
Maruyama, S., see Sasaki, S., 121
Marx, J., see Breidbach, O., 5
Matsui, T., see Nagafuji, T., 159
Matsumoto, M., see Kiuchi, Y., 193
McAdoo, D.J., see Paleckova, V., 1
McBride, P.A., see Jeffrey, M., 106
McCloskey, T.C., see Kehne, J.H., 101
McCulloch, J., see Park, C.K., 41
McGill, I.S., see Lantos, P.L., 67
McNeill, D.L., see Papka, R.E., 224
McNeill, D.L., Traugh Jr., N.E., Vaidya, A.M., Hua, H.T. and Papka, R.E., Origin and distribution of NADPH-diaphorase-positive neurons and fibers innervating the urinary bladder of the rat, 33
Melino, G., see Corasaniti, M.T., 221
Mongini, T., Palmucci, L., Doriguzzi, C., Chiadò-Piat, L. and Restagno, G., Absence of dystrophin in two patients with Becker type Xp21 muscular dystrophy, 37
Moroji, T., see Yamamoto, H., 97
Murakami, F., see Saito, Y., 81
Müsch-Nittel, K., see Bloms, P., 155
Müsch-Nittel, K., see Mußhoff, U., 163
Mußhoff, U., Madeja, M., Bloms, P., Müsch-Nittel, K. and Speckmann, E.-J., Tunicamycin-induced inhibition of functional expression of glutamate receptors in *Xenopus* oocytes, 163
Mußhoff, U., see Bloms, P., 155

Nagafuji, T., Matsui, T., Koide, T. and Asano, T., Blockade of nitric oxide formation by N^{ω}-nitro-L-arginine mitigates ischemic brain edema and subsequent cerebral infarction in rats, 159
Nagatomo, T., see Inenaga, K., 189
Nagatsu, I., see Komori, K., 197
Nicoll, R.A., see Solís, J.M., 16
Nisticò, G., see Corasaniti, M.T., 221
Nomura, M., see Tanaka, J., 111
Nörenberg, W., Gebicke-Haerter, P.J. and Illes, P., Inflammatory stimuli induce a new K^+ outward current in cultured rat microglia, 171
Nouillot, P., Bouisset, S. and Do, M.C., Do fast voluntary movements necessitate anticipatory postural adjustments even if equilibrium is unstable?, 1

Oguchi, K., see Kiuchi, Y., 193
Okawa, K., see Saito, Y., 81
Okuwa, M., see Yamamoto, H., 97
Olverman, H.J., see Griffiths, R., 175
Oostveen, J.A., see Williams, L.R., 136
Osherov, A., Goldberg, D. and Atlas, D., Is the neuronal voltage-sensitive calcium channel of human brain, human pancreatic β cells and rat brain much larger than predicted by its cDNA sequence?, 233
Otten, U., see Laudiero, L.B., 9

Palecek, J., see Paleckova, V., 1
Paleckova, V., Palecek, J., McAdoo, D.J. and Willis, W.D., The non-NMDA antagonist CNQX prevents release of amino acids into the rat spinal cord dorsal horn evoked by sciatic nerve stimulation, 1
Palfreyman, M.G., see Kehne, J.H., 101
Palmucci, L., see Mongini, T., 37
Papka, R.E., see McNeill, D.L., 33
Papka, R.E. and McNeill, D.L., Distribution of NADPH-diaphorase-positive nerves in the uterine cervix and neurons in dorsal root and paracervical ganglia of the female rat, 224
Park, C.K., McCulloch, J., Kang, J.K. and Choi, C.R., Efficacy of D-CPPene, a competitive *N*-methyl-D-aspartate antagonist in focal cerebral ischemia in the rat, 41
Park, D.H., see Kang, U.J., 53
Pawelzik, H., Dodt, H.-U. and Zieglgänsberger, W., Actions of vasoactive intestinal polypeptide (VIP) on neocortical neurons of the rat in vitro, 167

Raos, V., see Chen, S., 85
Restagno, G., see Mongini, T., 37
Roberts, G.W., see Lantos, P.L., 67
Rogawski, M.A., see Subramaniam, S., 213
Rossor, M.N., see Lantos, P.L., 67

Sagi, N., see Yamamoto, H., 97
Saito, Y., Murakami, F., Song, W.-J., Okawa, K., Shimono, K. and Katsumaru, H., Developing corticorubral axons of the cat form synapses on filopodial dendritic protrusions, 81
Sasaki, S. and Maruyama, S., Ultrastructural study of skein-like inclusions in anterior horn neurons of patients with motor neuron disease, 121
Schilter, D., see Laudiero, L.B., 9
Schünzel, G., see Wolf, G., 63
Schwarcz, R., see Du, F., 185
Scott, J.R., see Jeffrey, M., 106
Senyah, Y., see Kehne, J.H., 101
Seylaz, J., see Héron, A., 205
Sheldon, R.A., see Ferriero, D.M., 29
Shimono, K., see Saito, Y., 81
Singh, E.A., see Singh, V.K., 25
Singh, V.K., Yang, Y.-Y. and Singh, E.A., Immunoblot detection of antibodies to myelin basic protein in Alzheimer's disease patients, 25
Smith, C.B., see Lyon, M.J., 72
Sokoloff, L., see Lyon, M.J., 72
Solís, J.M. and Nicoll, R.A., Postsynaptic action of endogenous GABA released by nipecotic acid in the hippocampus, 16
Song, W.-J., see Saito, Y., 81
Speckmann, E.-J., see Mußhoff, U., 163
Speckmann, E.-J., see Bloms, P., 155
Spener, F., see Bloms, P., 155
Stroemer, R.P., Kent, T.A. and Hulsebosch, C.E., Increase in synaptophysin immunoreactivity following cortical infarction, 21
Subramaniam, S., Donevan, S.D. and Rogawski, M.A., 1,10-Diaminodecane and 1,12-diaminododecane block NMDA receptor currents by an open channel mechanism, 213
Swanson, R.A., Astrocyte glutamate uptake during chemical hypoxia in vitro, 143

Tabb, T., see Khan, I., 115
Tanaka, J., Hori, K., Koito, H. and Nomura, M., Enhanced monoamine release in the median preoptic area following reduced extracellular fluid volume in rats, 111
Tartaglia, R.L., see Corasaniti, M.T., 221
Ter-Margarian, A. and Djamgoz, M.B.A., Cytochalasin inhibits light-dependent synaptic plasticity of horizontal cells in teleost retina, 131
Traugh Jr., N.E., see McNeill, D.L., 33
Tseng, L.F., see Dun, N.J., 217

Ueta, Y., see Inenaga, K., 189

Vaidya, A.M., see McNeill, D.L., 33
Vanderhaeghen, J.-J., see Mailleux, P., 179, 182
Velluti, J.C., see Fortin, G., 89

Wagner, H.-J., see Behrens, U.D., 93
Walker, D.W., Lee, N., Heaton, M.B., King, M.A. and Hunter, B.E., Chronic ethanol consumption reduces the neurotrophic activity in rat hippocampus, 77
Wegerhoff, R., see Breidbach, O., 5
Wessel, T., see Kang, U.J., 53
Whatley, S.A., see Lantos, P.L., 67
Wiegandt, H., see Breidbach, O., 5
Williams, L.R. and Oostveen, J.A., Sensitivity of Fischer 344 × Brown Norway hybrid rats to exogenous NGF: weight loss correlates with stimulation of striatal choline acetyltransferase, 136
Willis, W.D., see Paleckova, V., 1
Wolf, G., Würdig, S. and Schünzel, G., Nitric oxide synthase in rat brain is predominantly located at neuronal endoplasmic reticulum: an electron microscopic demonstration of NADPH-diaphorase activity, 63
Woodgett, J.R., see Hanger, D.P., 58
Würdig, S., see Wolf, G., 63

Yamamoto, H., Sagi, N., Yamamoto, T., Goji, Y., Okuwa, M., Yoshii, M. and Moroji, T., Inhibitory effects of psychotomimetic σ ligands on nicotine-induced K^+ flux from differentiated PC12 cells, 97
Yamamoto, T., see Yamamoto, H., 97
Yamaoka, K., see Komori, K., 197
Yamashita, H., see Inenaga, K., 189
Yamazaki, Y., see Kiuchi, Y., 193
Yang, Y.-Y., see Singh, V.K., 25
Yoshii, M., see Yamamoto, H., 97

Zhang, F., see Mailleux, P., 182
Zieglgänsberger, W., see Pawelzik, H., 167